

UREDINALES COLLECTED BY FRED J. SEAVER IN TRINIDAD*

J. C. ARTHUR

No attempt has heretofore been made to enumerate the rusts of Trinidad. The present list of 71 species makes a creditable beginning. Probably two or three times this number may be found when a more thorough exploration is made. Over 160 species are already known for the island of Porto Rico, and about the same number for Cuba. Porto Rico has not one tenth the area of Cuba, and Trinidad has not quite one half the area of Porto Rico, but has a diversified topography and nearness to the mainland that will doubtless largely compensate for lessened area.

The island of Trinidad lies so close to the coast of South America that it is more properly considered a part of the southern continent rather than of the northern. Its flora is excluded from the volumes of the North American Flora.

The collections made by Dr. Seaver were obtained during the six weeks between March 1 and April 14, 1921. The visit to the island was made in the company of Dr. N. L. Britton,¹ who supplied a preliminary determination of the hosts in the field. The hosts have since been checked over by Mr. Percy Wilson at the herbarium of the New York Botanical Garden.

It is worthy of note that although Dr. Seaver gave his chief attention to other groups of fungi, yet he was able to secure 169 collections of Uredinales, which have yielded 71 species of rusts, 3 being new, or 4 including the one supplied by Mr. Nowell, as well as quite a number that are little known.

In the following list space has been economized by omitting the exact localities, except for new species, and by referring to the page of the seventh volume of the North American Flora, where

* Contribution from the Botanical Department of Purdue University Agricultural Experiment Station.

¹ For a detailed account of this trip, see Journal of the New York Botanical Garden for May, 1921.

the synonymy can be found, so far as that work has been published. All collections, except the one noted, are to be credited to Dr. Seaver, and the date of collection is March–April, 1921. Numbers 26, 33, 39, 41, 46, 47, 50, 52–57, thirteen in all, or eighteen per cent, are short-cycle species, the remaining fifty-eight are various forms of long-cycle species.

1. COLEOSPORIUM IPOMOEAE (Schw.) Burr. N. A. F. 87

On *Ipomoea glabra* (Aubl.) Choisy, II, 3181, 3275, 3291; *Ipomoea* sp., II, 3274, 3385. A common rust in its uredinal stage throughout tropical America.

2. PHAKOPSORA CROTONIS (Cooke) Arth. N. A. F. 104

On *Croton gossypifolium* L., II, 3253, 3424; *C. hirtus* L'Her., II, 3109. This rust was described by P. Hennings (*Hedwigia* 35: 251. 1896) under the name *Uredo crotonicola*, on *Croton grandulosus* from Argentina.

3. PHAKOPSORA MEIBOMIAE Arth. Bull. Torrey Club 44: 509.

1917

On *Meibomia supina* (Sw.) Britton, II, 3197; *M. triflora* (L.) Kuntze, II, 2960.

4. CEROTELIUM GOSSYPII (Lagerh.) Arth. N. A. F. 187

On *Gossypium* sp., II, 2953, 3388.

5. RAVENELIA INDIGOFERAE Tranz. N. A. F. 144

On *Indigofera suffruticosa* Mill. (*I. Anil* L.), 3077.

6. PROSPODIUM APPENDICULATUM (Wint.) Arth. N. A. F. 160

On *Tecoma Stans* (L.) Juss., III, 3406, 3430.

7. *Prosopidium suppressum* sp. nov.

O. Pycnia amphigenous, numerous, blackish-brown, inconspicuous, subcuticular, conic, small, about 100 μ broad and about same height; ostiolar filaments wanting.

II. Uredinia at first (primary) amphigenous, interspersed with the pycnia in circular groups 2-4 mm. across on somewhat larger discolored spots, afterward (secondary) hypophyllous, scattered, minute, 0.1-0.2 mm. in diameter, soon naked, pulverulent, chestnut-brown; paraphyses few or wanting; urediniospores flattened laterally, seen from flattened side globoid or somewhat obovoid, 23-32 by 25-40 μ , seen from narrow side obovate, and apparently acute above; wall with a hygroscopic layer, the inner layer firm, cinnamon-brown, 3 μ thick, the cuticle and subcuticular layers colorless, verrucose-echinulate with close-set, blunt projections on the flattened sides, becoming much longer on the narrowed sides and, as the spores are ordinarily⁹ seen, appearing like a coarse fringe, 5-7 μ wide, extending around the spore, the pores 2, distinct, one in center of each flattened side.

III. Telia hypophyllous, in loose groups, at first bullate, soon naked, somewhat pulverulent, chocolate-brown, ruptured cuticle noticeable; teliospores ellipsoid, 21-24 by 25-34 μ , rounded above and below, slightly constricted at septum; wall dark chestnut-brown, uniformly 2-3 μ thick, sparingly and evenly verrucose; pedicel colorless, as long as spore, slender, tapering downward, the appendages obsolete.

On *Tabebuia* sp. (Bignoniaceae), Las Lilas, March 24-28, O, II, 3350 (type); Pointe Gourde, March 31, II, III, 3408; Forest, Siparia Quarry, April 8, III, 3526; Lady Chancellor Road, March 14, II, iii, 3180. The group of species to which this new species belongs is imperfectly understood. There are probably quite a number of them, but at present the collections are frequently unnamed or listed under the genus *Uredo* or *Puccinia*, and have not been brought together for comparative study.

8. *UROMYCES AFFINIS* Wint. Hedwigia 24: 259. 1885

On *Hypoxis decumbens* L., II, 3192, 3199, 3387. The aecia and telia of this species were found in Missouri in 1883 by Demetrio, the earliest collection of the rust known, but so few urediniospores were present that they were not mentioned in the original description. Since that time many collections showing abundance of uredinia have been taken in the eastern United States, and recently the tropical collections, which show only uredinia and are usually reported under the names *Uredo Hypoxidis* (Bres.) P. Henn. and *Uredo globulosa* Arth., have been referred to the same species.

9. UROMYCES APPENDICULATUS (Pers.) Fries, N. A. F. 257
On *Phaseolus* sp., 3249.
10. UROMYCES BIDENTICOLA (P. Henn.) Arth. Mycologia 9: 71.
1917
On *Bidens pilosa* L., II, 3075, 3190.
11. UROMYCES COLUMBIANUS Mayor, Mém. Soc. Neuch. 5: 467.
1913
On *Melanthera aspera* (Jacq.) Steud., I, II, 3245; II, 3239,
3301.
12. UROMYCES DOLICHOLI Arth. N. A. F. 258
On *Cajan Cajan* (L.) Millsp., 3028, 3335, 3487.
13. UROMYCES HEDYSARI-PANICULATI (Schw.) Parl. N. A. F. 248
On *Meibomia Scorpiurus* (Sw.) Kuntze, 3194; *M. affinis* (Sch.)
Kuntze, 3306.
14. UROMYCES HOWEI Peck, N. A. F. 264
On *Asclepias Curassavica* L., II, 3304.
15. UROMYCES JANĪPHAE (Wint.) Arth. Mycologia 7: 190. 1915
On *Manihot Manihot* (L.) Cockerell, II, 3268, 3279, 3423.
Uredinia of this rust are common in tropical America, but the
aecia and telia have only been found in Mexico.
16. UROMYCES LEPTODERMUS Sydow, N. A. F. 224
On *Lasaicis* sp., II, 3402, 3477.
17. UROMYCES MAJOR Arth. (*Uredo ignobilis* Arth.), N. A. F. 225
On *Sporobolus indicus* (L.) R. Br., II, 3093.
18. UROMYCES NEUROCARPI Diet. N. A. F. 258
On *Clitoria rubiginosa* Juss., II, III, 3206, 3292.
19. UROMYCES PROËMINENS (DC.) Pass., N. A. F. 259
On *Chamaesyce hirta* (L.) Millsp., 2965, 2973, 3302, 3436, 3528.

20. UROMYCES SCLERIAE P. Henn. N. A. F. 233

On *Scleria melaleuca* Schlecht. & Cham., II, 3299. It has been reported from Cuba, Porto Rico, and southern Brazil.

21. UROMYCES WULFFIAE-STENOGLOSSAE Diet. Ann. Myc. 6: 96.
1908

On *Wulffia baccata* (L. f.) Kuntze, I, 2971; II, iii, 3244; I, II, iii, 3231.

22. PUCCINIA ACNISTI Arth. N. A. F. 471

On *Acnistus arborescens* Schlecht., I, 3178, 3227, 3524.

23. PUCCINIA AEQUINOCTIALIS Holway, Ann. Myc. 3: 22. 1905

On *Cydista aequinoctialis* (L.) Miers, II, 3092.

24. PUCCINIA ANGUSTATOIDES R. E. Stone, N. A. F. 351

On *Rynchospora cyperoides* (Sw.) Mart., III, 2976.

25. PUCCINIA ANTIOQUIENSIS Mayor, N. A. F. 347

On *Cyperus diffusus* Vahl, II, iii, 3337, 3393; III, 3269. The three collections show many 1-celled teliospores (mesospores), and the few teliospores found on 3337 and 3393 were all 1-celled, which would entitle these two collections to be entered under the genus *Uromyces*. The number of teliospores present are too few, however, to warrant the introduction of a new name.

26. PUCCINIA ARECHAVALETAE Speg. An. Soc. Ci. Arg. 12: 67.
1881

On *Urvillea Seriana* (L.) H. B. K., 2966, 3133, 3139, 3345, 3479.

27. PUCCINIA BIGNONIACEARUM Speg. An. Soc. Ci. Arg. 26: 11.
1888

On *Bignoniaceae*, III, 3333, 3378.

28. PUCCINIA CANNAE (Wint.) P. Henn. N. A. F. 380

On *Canna* sp., II, 3105, 3201, 3483; *Maranta arundinacea* L., II, 2969, 3091, 3138, 3141, 3476. The latter host has not before been reported.

29. PUCCINIA DEFORMATA Berk. & Curt. N. A. F. 294
On *Olyra latifolia* L., ii, III, 3474.
30. PUCCINIA EUPATORII-COLUMBIANI Mayor, Mém. Soc. Neuch.
5: 514. 1913
On *Eupatorium inulaefolium* H. B. K., II, III, 3587.
31. PUCCINIA GOUANIAE Holway, Ann. Myc. 3: 21. 1905
On *Gouania polygama* (Jacq.) Urban, II, 3027, 3457, 3090,
3254; II, III, 3478, 3117.
32. PUCCINIA HELICONIAE (Diet.) Arth. Bull. Torrey Club 45:
144. 1918
On *Heliconia psittacorum* (L. f.) Kuntze, II, 3202; *Heliconia*
sp., II, 3525.
33. PUCCINIA HETEROSPORA Berk. & Curt. Jour. Linn. Soc. 10:
356. 1868
On *Abutilon giganteum* (Jacq.) Presl, 3458.
34. PUCCINIA HYDROCOTYLES (Link) Cooke, Grevillea 9: 14.
1880
On *Hydrocotyle Hazeni* Rose, II, 3426.
35. PUCCINIA HYPTIDIS (M. A. Curt.) Tr. & Earle, N. A. F. 408
On *Hyptis capitata* (L.) Jacq., II, 2968.
36. PUCCINIA HYPTIDIS-MUTABILIS Mayor, N. A. F. 410
On *Hyptis mutabilis* (A. Rich.) Briq., I, 3074, 3386; II, 3107,
3189.
37. *Puccinia* (?) *ignava* comb. nov. (*Uredo ignava* Arth.), N.
A. F. 341
On *Bambos* sp., II, 2958, 3111. Although the teliospores of
this species are not known, it is highly probable that they will
eventually be found to conform to the requirements of the genus
Puccinia, and for convenience it is now so listed.

38. PUCCINIA IMPEDITA Mains & Holway; Arth. Mycologia 10:
135. 1918
On *Salvia occidentalis* Sw., II, 2962, 3272.
39. PUCCINIA LANTANAE Farl. Proc. Am. Acad. Sci. 18: 83.
1883
On *Priva lappulacea* (L.) Pers., 2955, 2970, 3397.
40. PUCCINIA LEONOTIDIS (P. Henn.) Arth. N. A. F. 407
On *Leonotis nepetaefolia* (L.) R. Br., II, 3273, 3354.
41. PUCCINIA OBLIQUA Berk. & Curt.; Berk. Jour. Linn. Soc. 10:
356. 1869
On *Metastelma* sp., 3188, 3448, 3488.
42. PUCCINIA PALLESCENS Arth. (*Uredo pallida* D. & H.),
N. A. F. 278
On *Zea Mays* L., II, 3103, 3110. No teliospores of this rust have yet been found on corn (maize), and the aecia are unknown.
43. PUCCINIA RUELLIAE (Berk. & Br.) Lagerh. N. A. F. 415
On *Blechnum Blechnum* (L.) Millsp. (*B. Brownei* Juss.), II, 2957, 3096, 3195; *Diantha pectoralis* (Jacq.) Gmel. (*Justicia pectoralis* Jacq.), II, 2954, 3191. The second host is a new record for the species.
44. PUCCINIA SCLERIAE (Paz.) Arth. (*Aecidium passifloricola*
P. Henn.) N. A. F. 349
On *Passiflora rubra* L., I, 3422.

45. **Puccinia Seaveriana** sp. nov.

II. Uredinia amphigenous, sparsely grouped or singly on yellowish spots, irregularly rounded, 0.1–0.5 mm. across, at first bul-
late, soon naked, somewhat pulverulent, cinnamon- or chestnut-
brown, ruptured epidermis conspicuous; paraphyses peripheral,
abundant, strongly incurved, cylindric, sometimes inclined toward
capitate, 10–15 by 40–50 μ , the wall thin, 1 μ , and pale or colorless
below, much thickened above, 3–6 μ , and dark chestnut-brown;
urediniospores broadly ellipsoid or globoid, 16–22 by 18–27 μ ; wall
cinnamon-brown, thin, 1–1.5 μ , closely and conspicuously echinu-
late, the pores 3, equatorial.

III. Telia not seen; teliospores in the uredinia oblong, 25–30 by 40–45 μ , rounded above and below, slightly or not constricted at septum; wall dark chestnut-brown, uniformly thick, about 3 μ , closely and noticeably verrucose; pedicel colorless, slender, fragile, once length of spore or shorter.

On *Oliganthes condensatus* (Less.) Schr. Bip. (*Carduaceae*), Lady Chancellor Road, March 14, II, 3179; same, March 17, II, 3236; same, March 21, II, 3283 (type); *Oliganthes Milleri*(?), western end of Monos Island, April 4, II, iii, 3459. The hosts belong to the tribe Vernonieae, and are part of a genus comprising about eight species confined to tropical America. All the species are trees or shrubs, *Oliganthes condensatus* producing the largest individuals known among the composites. The rust is notable for its abundance of deeply colored paraphyses. Such structures have been recorded for only one other species on the tribe Vernonieae. Only a few teliospores were found. The type for the species has been chosen to show the most characteristic and best development of the uredinia, although the presence of teliospores could not be demonstrated. The few teliospores seen were on a collection which also had but few uredinia. The specific name is selected to give recognition to the devotion of the collector of this and other material which is the basis of this report, by which he has added greatly to the store of mycological knowledge.

46. **Puccinia solanita** (Schw.) comb. nov. (*Aecidium solanitum* Schw. Jour. Acad. Sci. Phila. II, 2: 283. 1853; *Puccinia claviformis* Lagerh. Tromsö Mus. Aarsh. 17: 53. 1895)

On *Solanum* sp., 3295. The type collection for *Aecidium solanitum* is amply represented in the Schweinitz herbarium at the Philadelphia Academy of Science. An examination of this material shows that it bears a short-cycle rust, identical in appearance with *Puccinia claviformis*. The collection has also been examined by Mr. Percy Wilson and Dr. J. K. Small of the New York Botanical Garden, and they pronounce the host to be a species of *Solanum*, possibly *S. Melongena* L.

47. Puccinia SPEGAZZINII DeToni in Sacc. Syll. 7: 704. 1888

On *Mikania micrantha* H. B. K., 3527; *Mikania* sp., 2956, 3135, 3228, 3312.

48. PUCCINIA STRIOLATA (Speg.) Arth. (*P. macropoda* Speg.)
N. A. F. 387
On *Iresine Celosia* L., I, II, 3137.
49. PUCCINIA SUBSTRIATA Ell. & Barth. N. A. F. 289
On *Eriochloa punctata* Ham., II, 3193.
50. PUCCINIA SYNEDRELLAE P. Henn. Hedwigia 37: 277. 1898
On *Synedrella nodiflora* (L.) Gaertn., 2964, 3271, 3586; *Emilia sonchifolia* (L.) DC., 3073.
51. PUCCINIA TUBULOSA (Pat. & Gaill.) Arth. (*Uredo paspalicola* P. Henn.) N. A. F. 288
On *Paspalum paniculatum* L., ii, 3112; *Paspalum* sp., II, 2961, 3185; *Syntherisma digitata* Hitchc., II, 3339. The aecia of this species occur on *Solanum torvum* and closely related hosts.
52. PUCCINIA URBANIANA P. Henn. Hedwigia 37: 278. 1898
On *Valerianodes cayennensis* (Vahl) Kuntze, 2967, 2972, 3255.
53. ENDOPHYLLUM CIRCUMSCRIPTUM (Schw.) Whetzel & Olive,
Am. Jour. Bot. 4: 49. 1917
On *Cissus sicyoides* L., 3267.
54. ENDOPHYLLUM DECOLORATUM (Schw.) Whetzel & Olive, l. c.
On *Clibadium surinamense* L., 3177, 3270.
55. ENDOPHYLLUM WEDELIAE (Earle) Whetzel & Olive, l. c.
On *Wedelia trilobata* (L.) Hitchc., 3087.
56. ENDOPHYLLOIDES PORTORICENSIS Whetzel & Olive, l. c.
On *Mikania* sp., 3136, 3523.
57. PUCCINIOSIRA PALLIDULA (Speg.) Lagerh. N. A. F. 127
On *Triumfetta* sp., 3079, 3248b, 3252, 3310.
58. UREDO ADENOCALYMMATIS P. Henn. Hedwigia 35: 249. 1896
On Bignoniaceae, 3203, 3340. This rust, both in its spores and paraphyses, has a close resemblance to species which have been

referred to the genus *Prospodium*. The paraphyses are notable for their scimitar shape, sharp points, and cross walls. It has been reported on *Pyrostegia venusta*, as well as on the type genus.

59. UREDO COMMELYNAE Kalchbr. *Grevillea* 11: 24. 1882
On *Commelina elegans* H. B. K., 3390.

60. UREDO CYATHULAE Mayor, *Mém. Soc. Neuch.* 5: 584. 1913
On *Cyathula achyranthoides* (H. B. K.) Moq., 3334.

61. UREDO MACULANS Pat. & Gaill. *Bull. Soc. Myc. Fr.* 4: 98.
1888

On *Pfaffia iresinoides* (H. B. K.) Kuntze, 3076, 3250, 3303.

62. UREDO MANDEVILLAE Mayor, *Mém. Soc. Neuch.* 5: 591. 1913
On *Mandevilla tomentosa* (Vahl) Kuntze, 3258, 3285, 3297.

63. UREDO RUBESCENS Arth. *Mycologia* 7: 327. 1915
On *Dorstenia Contrajerva* L., 3078, 3475.

64. UREDO TRICHILIAE Arth. *Mycologia* 9: 90. 1917
On *Trichilia trinitensis* A. Juss., 3305, 3421.

65. UREDO VICINA Arth. *Mycologia* 7: 325. 1915
On *Wedelia Jacquinii* Rich., 3432.

66. UREDO VITICIS Juel, *Bih. K. Sv. Vet.-Akad. Handl.* 23(3)¹⁰:
26. 1897

On *Vitex* sp., 3293.

67. *Aecidium Alibertiae* sp. nov.

O. Pycnia epiphyllous, numerous in circular groups on discolored spots 4–10 mm. across, prominent, subepidermal but appearing subcuticular, flattened-conic, large, 160–210 μ in diameter, 65–80 μ high; hymenium flat; ostiolar filaments wanting.

I. Aecia hypophyllous, opposite the pycnia, short-cylindric, 0.3–0.4 mm. in diameter, deep-seated, extending half way through the leaf; peridium colorless, the margin coarsely lacerate, fragile; peridial cells in front view angularly ellipsoid or oblong, in side view lanceolate, strongly overlapping, 16–20 by 30–40 μ , the outer wall thin, 1–2 μ , smooth, the inner wall thicker, 2–5 μ , moderately

and closely verrucose; aeciospores globoid, 21-26 by 23-29 μ ; wall pale or cinnamon-brown, 1.5-2 μ thick, finely and closely verrucose.

On *Alibertia* sp. (Rubiaceae), Piarco savanna, March 15, 3204; Piarco savanna, south of Dabadie, March 21, 3286 (type); Meara savanna, March 22, 3296. The species is remarkable for the large pycnia, that are formed beneath the thin epidermis, but above the thick palisade cells. They are morphologically similar to subcuticular pycnia.

68. *AECIDIUM BRASILIENSE* Diet. Hedwigia 36: 35. 1897

On *Cordia cylindrostachya* R. & S., 3106, 3246, 3251, 3277.

69. *AECIDIUM BYRSONIMATIS* P. Henn. Hedwigia 34: 101. 1895

On *Byrsonima verbascifolia* Rich. (?), 3200.

70. *Aecidium delicatum* sp. nov.

O. Pycnia amphigenous, in small close groups, punctiform, honey-yellow, noticeable, subepidermal, globoid, about 125 μ in diameter.

I. Aecia hypophyllous, surrounding the pycnia, somewhat circinating, on yellowish spots 1-2 mm. across, low and broad, 0.5-0.8 mm. in diameter; peridium delicate, erect, finely erose; peridial cells oblong in surface view, rhomboidal in side view, slightly overlapping, 26-32 μ long, the wall colorless, the inner wall 3-5 μ thick, finely verrucose, the outer wall thinner, smooth; aeciospores globoid or ellipsoid, 16-24 by 20-30 μ ; wall colorless, thin, 1 μ , minutely and closely verrucose.

On *Eucharis* sp. (Amaryllidaceae), Port of Spain, no date, collected by Nowell and communicated by Seaver. Little comparative study has been made of the rusts on Amaryllidaceous hosts. Their identity is made especially difficult by the collection of single stages and on hosts not fully determined.

71. *AECIDIUM TOURNEFORTIAE* P. Henn. Hedwigia 34: 338.

1895

On *Tournefortia tomentosa* Mill., 3278.

INDEX TO UREDINALES

(New or newly combined names are in **bold face type**)

- Aecidium* **Alibertiae** 67
brasiliense 68
Byrsonimatis 69
delicatum 70
passifloricola 44
solanium 46
Tournefortiae 71
Cerotelium *Gossypii* 4
Coleosporium *Ipomoeae* 1
Endophylloides *portoricensis* 55
Endophyllum *circumscriptum* 52
decoloratum 53
Wedeliae 54
Phakopsora *Meibomia* 3
Prospodium *appendiculatum* 6
suppressum 7
Puccinia *Acnisti* 22
aequinoctialis 23
angustatoides 24
antioquiensis 25
Arechavaletae 26
Bignoniacearum 27
Cannae 28
claviformis 46
deformata 29
Eupatorii-columbiani 30
Gouaniae 31
Heliconiae 32
heterospora 33
Hydrocotyles 34
Hyptidis 35
Hyptidis-mutabilis 36
ignava 37
impedita 38
Lantanae 39
Leonotidis 40
macropoda 48
obliqua 41
pallescens 42
Ruelliae 43
Scleriae 44
Seaveriana 45
solanita 46
Spegazzinii 47
striolata 48
substriata 49
Synedrellae 50
tubulosa 51
Urbaniana 52
Puccinosira *pallidula* 77
Ravenelia *Indigoferae* 5
Uredo *Adenocalymmatiss* 58
Commelynae 59
crotonicola 2
Cyathulae 60
globulosa 8
Hypoxidis 8
ignava 37
ignobilis 17
maculans 61
Mandevillae 62
paspalicola 51
rubescens 63
Trichiliae 64
vicina 65
Viticis 66
Uromyces *affinis* 8
appendiculatus 9
bidenticola 10
columbianus 11
Dolicholi 12
Hedysari-paniculati 13
Howei 14
Janiphae 15
leptodermus 16
major 17
Neurocarpi 18
proëminens 19
Wulffiae-stenoglossae 21

HOST INDEX

- Abutilon giganteum* 33
Acnistus arborescens 22
Alibertia sp. 67
Asclepias Curassavica 14
Bambos sp. 37
Bidens pilosa 10
 Bignoniaceae 7, 27, 58
Blechnum Blechnum 43
 Brownei 43
Byrsonima verbascifolia 69
Cajan *Cajan* 12
Canna sp. 28
Chamaesyce hirta 19
Cissus sicyoides 53
Clibadium surinamense 54
Clitoria rubiginosa 18
Commelina elegans 59
Cordia cylindrostachya 68
Croton glandulosus 2
 gossypifolium 2
 hirtus 2
Cyathula achyranthoides 60
Cydista aequinoctialis 23
Cyperus diffusus 25
Dorstenia Contrajerva 63
Emilia sonchifolia 50
Eriochloa punctata 49
Eucharis sp. 70
Eupatorium inulaefolium 30
Gossypium sp. 4
Gouania polygama 31
Hydrocotyle Hazeni 34
Hypoxis decumbens 8
Hyptis capitata 35
 mutabilis 36
Indigofera suffruticosa 5
Ipomoea glabra 1
Iresine Celosia 48
 Lasaisis sp. 16
Leonotis nepetaefolia 40
Mandevilla tomentosa 62
Manihot Manihot 15
Maranta arundinacea 28
Meibomia affinis 13
 Scorpiurus 13
 supina 3
Melanthera aspera 11
Metastelma sp. 41
Mikania micrantha 47
 sp. 47, 56
Oliganthes condensatus 45
 Milleri 45
Olyra latifolia 29
Paspalum paniculatum 51
Passiflora rubra 44
Pfaffia iresinoides 61
Phaseolus sp. 9
Priva lappulacea 39
Rynchospora cyperoides 24
Salvia occidentalis 38
Solanum sp. 46
Sporobolus indicus 17
Synedrella nodiflora 50
Syntherisma digitata 51
Tabebuia sp. 7
Tecoma Stans 6
Tournefortia tomentosa 71
Trichilia trinitensis 64
Triumfetta sp. 57
Urvillea Seriana 26
Valerianodes cayennensis 52
Vitex sp. 66
Wedelia Jacquini 65
 trilobata 55
Wulffia baccata 21
Zea Mays 42

PURDUE UNIVERSITY,
 LAFAYETTE, INDIANA.